

Alert	<p>Imipenem + cilastatin is not the preferred carbapenem in neonates because of possible adverse effects and should be avoided in preterm neonates because of cilastatin accumulation.</p> <p>The Antimicrobial Stewardship Team recommends this drug is listed under the following category: Restricted.</p> <p>Widespread use of carbapenems has been linked with increasing prevalence of infections caused by methicillin-resistant <i>Staphylococcus aureus</i> (MRSA), vancomycin-resistant enterococci (VRE), multi resistant Gram-negative organisms and <i>Clostridium difficile</i>.</p>												
Indication	<p>Non-CNS sepsis caused by susceptible organisms including enteric Gram-negative rods, extended-spectrum beta-lactamase [ESBL] organisms, <i>Pseudomonas aeruginosa</i>, anaerobic organisms (including <i>Bacteroides fragilis</i>) and many Gram-positive organisms.</p>												
Action	<p>Imipenem + cilastatin is a carbapenem. It inhibits cell wall synthesis. Imipenem is combined with cilastatin. Cilastatin prevents renal metabolism of imipenem.</p> <p>Meropenem is a better choice than imipenem + cilastatin for central nervous system infections. Meropenem attains a higher concentration in the cerebrospinal fluid and has a lower incidence of seizures than imipenem + cilastatin.</p>												
Drug Type	<p>Carbapenem antibiotic.</p>												
Trade Name	<p>Primaxin</p>												
Presentation	<p>500 mg vial.</p>												
Dosage / Interval	<table border="1"> <thead> <tr> <th data-bbox="408 994 683 1028">Condition</th> <th data-bbox="683 994 951 1028">Dose</th> <th data-bbox="951 994 1219 1028">Dosing Interval</th> <th data-bbox="1219 994 1477 1028">Infusion Time</th> </tr> </thead> <tbody> <tr> <td data-bbox="408 1028 683 1095">Non-<i>Pseudomonas aeruginosa</i></td> <td data-bbox="683 1028 951 1095">25 mg/kg</td> <td data-bbox="951 1028 1219 1095">12 hourly</td> <td data-bbox="1219 1028 1477 1095">30 minutes</td> </tr> <tr> <td data-bbox="408 1095 683 1162"><i>Pseudomonas aeruginosa</i></td> <td data-bbox="683 1095 951 1162">25 mg/kg</td> <td data-bbox="951 1095 1219 1162">8 hourly</td> <td data-bbox="1219 1095 1477 1162">90 minutes</td> </tr> </tbody> </table>	Condition	Dose	Dosing Interval	Infusion Time	Non- <i>Pseudomonas aeruginosa</i>	25 mg/kg	12 hourly	30 minutes	<i>Pseudomonas aeruginosa</i>	25 mg/kg	8 hourly	90 minutes
Condition	Dose	Dosing Interval	Infusion Time										
Non- <i>Pseudomonas aeruginosa</i>	25 mg/kg	12 hourly	30 minutes										
<i>Pseudomonas aeruginosa</i>	25 mg/kg	8 hourly	90 minutes										
Route	<p>IV Infusion.</p>												
Maximum Daily Dose	<p>75 mg/kg/day</p>												
Preparation/Dilution	<p>Add 9.2 mL of sodium chloride 0.9% to the 500 mg powder for reconstitution to make a volume of 10 mL with a concentration of 50 mg/mL (Note: Suspension maybe cloudy).</p> <p>Draw up 1 mL (50 mg) and add 9 mL sodium chloride 0.9% to make a final volume of 10 mL with a concentration of 5 mg/mL.</p>												
Administration	<p>Non-<i>Pseudomonas aeruginosa</i> – IV infusion over 30 minutes. <i>Pseudomonas aeruginosa</i> – IV infusion over 90 minutes.</p>												
Monitoring	<p>Monitor renal function. Dose may need to be reduced in impaired renal function. Monitor blood count and liver function.</p>												
Contraindications	<p>Hypersensitivity to penicillins, cephalosporins or carbapenems. CNS infections.</p>												
Precautions	<p>Seizures can occur in infants with renal impairment or central nervous system infection.</p>												
Drug Interactions	<p>Ganciclovir – risk of seizures. Do not give concomitantly unless the potential benefits outweigh the risks. Valproate – results in decreased concentrations of valproate.</p>												
Adverse Reactions	<p>Seizures, impaired renal function, impaired liver function, tachycardia, local phlebitis. Urticaria, diarrhoea, pseudomembranous colitis (<i>Clostridium difficile</i>) and vomiting.</p>												

Compatibility	<p>Fluids: Glucose 5%, glucose 10%, sodium chloride 0.9%</p> <p>Y-site: Aciclovir, amifostine, anidulafungin, aztreonam, caspofungin, cisatracurium besilate, foscarnet, granisetron, linezolid, remifentanyl, tigecycline, zidovudine.</p>
Incompatibility	<p>Fluids: Hartmann's.</p> <p>Y-site: Amiodarone, amoxicillin, azathioprine, azithromycin, ceftriaxone, chlorpromazine, daptomycin, fluconazole, ganciclovir, haloperidol lactate, metaraminol, midazolam, milrinone, mycophenolate mofetil, palonosetron, pethidine, sodium bicarbonate, vecuronium.</p>
Stability	<p>Reconstituted or diluted solution: Stable for 4 hours below 25°C or for 24 hours at 2–8°C.</p>
Storage	<p>Vial: Store below 25°C.</p>
Special Comments	<p>Solutions of imipenem + cilastatin range from colourless to yellow. Variations of colour within this range do not affect the potency.</p>
Evidence summary	<p>Pharmacokinetics: Imipenem + cilastatin is excreted via kidneys, mainly through glomerular filtration. Imipenem clearance is not influenced by postnatal or postmenstrual age. Infusions (0.5 hours) of 25 mg/kg every 12 hours (50 mg/kg/day) is sufficient against common bacterial isolates in neonates. However, 1.5 hour infusions of 25 mg/kg every 8 hours (75 mg/kg/day) in neonates are required to be effective against <i>Pseudomonas aeruginosa</i>.¹</p> <p>Safety: Seizures can occur in neonates with meningitis, other CNS infections and in patients with renal impairment.^{1,4,6,9}</p>
References	<ol style="list-style-type: none"> 1. Yoshizawa K, Ikawa K, Ikeda K, Ohge H, Morikawa N. Population pharmacokinetic-pharmacodynamic target attainment analysis of imipenem plasma and urine data in neonates and children. <i>Pediatr Infect Dis J</i> [Internet]. 2013 [cited 2013 Nov];32(11):1208–16. http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=med1&NEWS=N&AN=23676856 2. Fujimura S, Nakano Y, Sato T, Shirahata K, Watanabe A. Relationship between the usage of carbapenem antibiotics and the incidence of imipenem-resistant <i>Pseudomonas aeruginosa</i>. <i>J Infect Chemother</i> [Internet]. 2007 [cited 2007 Jun];13(3):147–50. http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=med5&NEWS=N&AN=17593500 3. Schlossberg D, Pietroski N. Carbapenems. <i>Semin Pediatr Infect Dis</i> [Internet]. 2002 [cited 2002 Jan];13(1):4. http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=med4&NEWS=N&AN=12118842 4. Boswald M, Dobig C, Kandler C, Kruger C, Scharf J, Soergel F, Zink S, Guggenbichler JP. Pharmacokinetic and clinical evaluation of serious infections in premature and newborn infants under therapy with imipenem/cilastatin. <i>Infection</i> [Internet]. 1999 [cited 1999];27(4-5):299–304. http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=med4&NEWS=N&AN=10885853 5. Blumer JL. Pharmacokinetic determinants of carbapenem therapy in neonates and children. <i>Pediatr Infect Dis J</i> [Internet]. 1996 [cited 1996 Aug];15(8):733–7. http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=med4&NEWS=N&AN=8858691 6. Stuart RL, Turnidge J, Grayson ML. Safety of imipenem in neonates. <i>Pediatr Infect Dis J</i> [Internet]. 1995 [cited 1995 Sep];14(9):804–5. http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=med3&NEWS=N&AN=8559632 7. Reed MD, Kliegman RM, Yamashita TS, Myers CM, Blumer JL. Clinical pharmacology of imipenem and cilastatin in premature infants during the first week of life. <i>Antimicrob Agents Chemother</i> [Internet]. 1990 [cited 1990 Jun];34(6):1172–7. http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=med3&NEWS=N&AN=2393278 8. Ahonkhai VI, Cyhan GM, Wilson SE, Brown KR. Imipenem-cilastatin in pediatric patients: an overview of safety and efficacy in studies conducted in the United States. <i>Pediatr Infect Dis J</i> [Internet]. 1989 [cited 1989 Nov];8(11):740–4. http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=med3&NEWS=N&AN=2687787 9. Nalin DR, Jacobsen CA. Imipenem/cilastatin therapy for serious infections in neonates and infants. <i>Scand J Infect Dis Suppl</i> [Internet]. 1987 [cited 1987];5246–55. http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=med2&NEWS=N&AN=3331042

	<p>10. Micromedex 2.0 accessed via CIAP 4nd November 2015.</p> <p>11. Australian Injectable Drugs Handbook, 6th Edition, Society of Hospital Pharmacists of Australia 2015.</p> <p>12. Neofax accessed on www.neofax.micromedex.solutions.com on 16th December 2015.</p>
--	--

Original version Date: 05/12/2015	Author: NeoMed Consensus Group
Current Version number: 1	Version Date: 05/12/2015
Risk Rating: High	Due for Review: 05/12/2017
Approved by: As per Local policy	Approval Date: As per Local policy