

Alert	Risk of infantile hypertrophic pyloric stenosis is significantly higher in neonates treated with erythromycin. ¹⁶
Indication	<ol style="list-style-type: none"> 1. Pertussis – post-exposure prophylaxis and treatment (azithromycin is recommended). 2. Chlamydial conjunctivitis and pneumonia 3. Treatment of other susceptible bacterial infections in penicillin-allergic infants 4. Prokinetic agent for gastrointestinal dysmotility (routine use not recommended)
Action	Inhibits protein synthesis by attaching to the 50S subunit of the bacterial ribosome in susceptible organisms. Motilin receptor agonist.
Drug Type	Macrolide antibiotic.
Trade Name	E-Mycin Syrup, EES Granules
Presentation	200 mg/5 mL suspension (granules for reconstitution), 100 mL 400 mg/5 mL suspension (granules for reconstitution), 100 mL
Dosage / Interval	<p><u>Pertussis – post-exposure prophylaxis and treatment¹ (azithromycin is recommended. Use erythromycin only if azithromycin is not available).</u></p> <p>10 mg/kg/dose PO 6-hourly for 5 to 14 days; 14 days preferred to avoid risk of relapse.</p> <p><u>Chlamydia infection (conjunctivitis, pneumonia)</u></p> <p>12.5 mg/kg/dose PO 6-hourly for 14 days.²</p> <p><u>Non-chlamydial, susceptible bacterial infection in penicillin-allergic infants³</u></p> <p>Weight <1 kg: -Postnatal age ≤14 days: 10 mg/kg/dose 12-hourly -Postnatal age >14 days: 10 mg/kg/dose 8-hourly</p> <p>Weight ≥1 kg: -Postnatal age ≤7 days: 10 mg/kg/dose 12-hourly -Postnatal age >7 days: 10 mg/kg/dose 8-hourly</p> <p><u>Prokinetic dose for gastrointestinal dysmotility (inconsistent evidence for its efficacy and safety and routine use not recommended)</u></p> <p>Low-dose regimens: 2.5 mg/kg/dose 6-hourly up to 10 days.¹⁰ or 5 mg/kg/dose 8-hourly (7–14 days)¹¹ High dose regimens: Doses up to 10–12.5 mg/kg/dose 6-hourly for 7–14 days have been used.¹²⁻¹⁴ Post-op intestinal atresia: 3 mg/kg/dose 6-hourly¹⁹</p>
Route	Oral
Preparation/Dilution	Add 77 mL of sterile water to granules in small volumes and shake vigorously until no lumps are visible. Suspension expires 10 days after reconstitution. Write expiry date on bottle.
Administration	Oral, preferably with feeds. ¹⁵ For prokinetic effect, often administered 30 minutes prior to feed.
Monitoring	Liver function.
Contraindications	Hypersensitivity to erythromycin or any component of the product. Concomitant therapy with pimozide, cisapride, ergotamine or dihydroergotamine, terfenadine, astemizole, lovastatin or simvastatin.
Precautions	Use with caution in hepatic impairment. QT interval prolongation. Uncorrected hypokalaemia, hypomagnesaemia. Class 1A and Class 3 antiarrhythmic agents.
Drug Interactions	QT interval prolonging drugs: Cisapride, fluconazole, octreotide, cotrimoxazole, verapamil, Class 1A and Class 3 antiarrhythmic agents. Drugs that may increase toxicity of erythromycin: Ketoconazole. Drugs that may reduce erythromycin plasma concentration: Carbamazepine, theophylline. Erythromycin may increase plasma concentrations of following drugs: Carbamazepine, digoxin,

	theophylline, warfarin, midazolam.
Adverse Reactions	<p>Infantile hypertrophic pyloric stenosis (IHPS): Risk of developing IHPS following erythromycin exposure is 0.4 % (95% CI 0.3–0.5%) in those receiving erythromycin at any time and 2.6 % (95% CI 1.5–4.2%) in those receiving erythromycin in the first 14 days.¹⁶</p> <p>COMMON: Nausea, vomiting and abdominal pain. The incidence of GI reactions may vary with the erythromycin salt preparation and/or dosing regimen. Diarrhoea may occur due to increased gastrointestinal motility caused by erythromycin.</p> <p>LESS FREQUENT OR RARE: Pancreatitis, pyloric stenosis, ileus, pseudomembranous colitis, sensorineural hearing loss, cholestasis, acute hepatitis, hepatic failure, agranulocytosis, thrombocytopenia, haemolytic anaemia, hypothermia, hypovolaemic shock and hypotension, leukocytoclastic vasculitis, acute respiratory distress following an allergic reaction, Schonlein-Henoch syndrome, candidal esophagitis, gingival hyperplasia, contact dermatitis, fixed drug eruptions, toxic pustuloderma, toxic epidermal necrolysis, interstitial nephritis, glomerulonephritis.</p>
Compatibility	Not applicable
Incompatibility	Not applicable
Stability	After reconstituting granules, refrigerate and use within 10 days.
Storage	Store granules below 25°C. Reconstituted suspension should be refrigerated at 2–8°C and used within 10 days; do not freeze.
Special Comments	<p>Readily absorbed.</p> <p>Hepatic metabolism by cytochrome P450 enzymes.</p>
Evidence summary	<p>Efficacy</p> <p><u>Pertussis – post-exposure prophylaxis and treatment^{1,4}</u></p> <p>Systematic review of eradicating <i>B. pertussis</i> from the nasopharynx found short-term antibiotics (azithromycin for 3–5 days, or clarithromycin or erythromycin for 7 days) were as effective as long-term (erythromycin for 10 to 14 days) (risk ratio (RR) 1.01; 95% CI 0.98 to 1.04), but had fewer side effects (RR 0.66; 95% CI 0.52 to 0.83).⁴</p> <p>The Centers for Disease Control and Prevention recommends oral azithromycin 10 mg/kg/day daily for 5 days. Azithromycin has the advantage of once daily dosing and shorter duration of therapy.¹</p> <p>Erythromycin may be used if azithromycin is unavailable: 40 mg/kg per day in 4 divided doses for 14 days.¹</p> <p><u><i>Chlamydia</i> prophylaxis in infants born to mothers who have chlamydial infection²</u></p> <p>Infants born to mothers who have untreated chlamydia are at high risk for infection. However, prophylactic antibiotic treatment is not indicated and the efficacy of such treatment is unknown. Infants should be monitored to ensure appropriate diagnosis and treatment if symptoms develop.</p> <p><u>Treatment of chlamydial conjunctivitis and pneumonia²</u></p> <p><i>C. trachomatis</i> infection in neonates is most frequently recognised by conjunctivitis that develops 5 to 12 days after birth. <i>C. trachomatis</i> also can cause a subacute, afebrile pneumonia with onset at ages 1 to 3 months. RCTs reported chlamydial conjunctivitis or pneumonia is eradicated after systemic treatment with oral erythromycin 50 mg/kg/day for 14 days with few treatment failures and is more effective than topical treatment for chlamydia conjunctivitis.⁵⁻⁸</p> <p>Recommendation: The Centers for Disease Control and Prevention recommends oral erythromycin 50 mg/kg per day given orally in four divided doses for 14 days for either chlamydial conjunctivitis or pneumonia. Alternative regimen is azithromycin 20 mg/kg/day, once daily for 3 days. Topical antibiotic therapy alone is inadequate and is unnecessary when systemic treatment is administered.²</p> <p><u>Prokinetic agent in preterm infants</u></p> <p>Systematic review evaluated the efficacy of erythromycin for prophylaxis or treatment of feeding intolerance in preterm infants. Ng and Shah 2008⁹ reviewed 10 randomised, controlled studies using both high- and low-dose erythromycin. Meta-analysis on most outcomes couldn't be done. Erythromycin for prevention or treatment demonstrated no consistent effect on time required to achieve full feeds. Three studies using erythromycin at doses between 40 and 50 mg/kg/day</p>

	<p>reported a statistically significant effect on feeding tolerance as did one study using erythromycin at a slightly smaller dose (but still considered high dose) of 15 mg/kg/day. A single study (Oei 2001) using low-dose erythromycin (10 mg/kg/day) for prevention of feed intolerance reported showed that infants in the erythromycin group achieved full feeds significantly earlier than the placebo group.¹⁰ However, three other studies that used low-dose erythromycin failed to show any significant difference between erythromycin and placebo in the times to establish full feeds in preterm infants <32 weeks' gestation with feeding intolerance. There was no reported effect on other neonatal morbidities including necrotising enterocolitis or sepsis.</p> <p>Conclusion: Although some studies have reported a reduced time to full feeds, the effect is inconsistent, the optimal dose is unclear and there has been no reported consistent effect on other neonatal morbidities.⁸ (LOE I, GOR C)</p> <p><u>Prokinetic agent in surgical infants</u></p> <p>An RCT comparing erythromycin 3 mg/kg/dose 4 times daily compared with placebo after primary repair of uncomplicated gastroschisis in 62 infants reported no difference in time to achieve full enteral feeding (27.2 v 28.7 days; P = .75), catheter-related sepsis, duration of parenteral nutrition or time to discharge between the 2 groups. [1] An RCT comparing erythromycin 3 mg/kg/dose 4 times daily in 30 neonates undergoing primary anastomosis for congenital small bowel atresia reported neonates receiving oral erythromycin achieved full enteral feeding earlier (13.07 vs. 16.13 days), required PN for a shorter duration (10.53 vs. 13.73 days) and their hospital stay was less (16.2 vs. 18.0 days). [2] Conclusion: There is inconsistent evidence that erythromycin 3 mg/kg/dose 4 times daily may have a beneficial effect in newborn infants with abdominal surgical conditions restricted to infants undergoing repair of small intestinal atresia. (LOE II GOR D)^{18,19}</p> <p>Safety</p> <p>A systematic review of observational data reported an increase in the absolute risk of developing infantile hypertrophic pyloric stenosis (IHPS) following erythromycin exposure of 0.4 % (95% CI 0.3–0.5%) in those receiving erythromycin at any time, and 2.6% (95% CI 1.5–4.2%) in those receiving erythromycin in the first 14 days.¹⁶</p> <p>Bioavailability</p> <p>The absorption was lower in infants <1 month of age than in older children. Administration of the drug with feeds considerably increased the absorption of erythromycin ethylsuccinate.¹⁵</p>
References	<ol style="list-style-type: none"> Centers for Disease Control and Prevention. Recommended Antimicrobial Agents for the Treatment and Postexposure Prophylaxis of Pertussis, 2005. MMWR Recomm Rep. 2005 December 9, 2005 / 54(RR14);1-16. Centers for Disease Control and Prevention. Sexually transmitted diseases treatment guidelines, 2015. MMWR Recomm Rep. 2015 Jun 5;64(RR-03):1-137. American Academy of Pediatrics (AAP). In: Pickering LK, Baker CJ, Kimberlin DW, Long SS, eds. Red Book: 2012 Report of the Committee on Infectious Diseases. 29th ed. Elk Grove Village, IL: American Academy of Pediatrics; 2012. Altunajji SM, Kukuruzovic RH, Curtis NC, Massie J. Antibiotics for whooping cough (pertussis). Cochrane Database of Systematic Reviews. 2007. Heggie AD, Jaffe AC, Stuart LA, Thombre PS, Sorensen RU. Topical sulfacetamide vs oral erythromycin for neonatal chlamydial conjunctivitis. American Journal of Diseases of Children. 1985;139:564-6. Sandstrom I. Treatment of neonatal conjunctivitis. Arch Ophthalmol. 1987;105:925-8. Stenberg K, Mardh PA. Treatment of chlamydial conjunctivitis in newborns and adults with erythromycin and roxithromycin. J Antimicrob Chemother. 1991;28:301-7. Lee PI, Wu MH, Huang LM, Chen JM, Lee CY. An open, randomized, comparative study of clarithromycin and erythromycin in the treatment of children with community-acquired pneumonia. J Microbiol Immunol Infect. 2008;41:54-61. Ng E, Shah VS. Erythromycin for the prevention and treatment of feeding intolerance in preterm infants. Cochrane Database of Systematic Reviews 2008, Issue 3. Art. No.: CD001815. DOI:

	<p>10.1002/14651858.CD001815.pub2.</p> <p>10. Oei J, Lui K. A placebo-controlled trial of low-dose erythromycin to promote feed tolerance in preterm infants. <i>Acta Paediatrica</i>. 2001 Aug 1;90(8):904-8.</p> <p>11. Ng SC, Gomez JM, Rajadurai VS, Saw S, Quak S. Establishing enteral feeding in preterm infants with feeing intolerance; A randomized controlled study of low-dose erythromycin. <i>JPGN</i> 2003;37:554-8.</p> <p>12. Madani A, Pishva N, Pourarian Sh, ZarkeshM. The efficacy of oral erythromycin in enhancement of milk tolerance in premature infants: A randomized controlled trial. <i>Iranian Journal of Medical Sciences</i> 2004;29:1-4.</p> <p>13. Ng PC, So KW, Fung KSC, Lee CH, Fok TF, Wong E, et al. Randomised controlled study of oral erythromycin for treatment of gastrointestinal dysmotility in preterm infants. <i>Archives of Disease in Childhood. Fetal and Neonatal Edition</i> 2001;84:F177-82.</p> <p>14. Nuntnarumit P, Kiatchosakun P, Tantiprapa W, Boonkasidecha S. Efficacy of oral erythromycin for treatment of feeding intolerance in preterm infants. <i>Journal of Pediatrics</i> 2006;148:600-5.</p> <p>15. Eriksson M, Bolme P, Blennow M. Absorption of erythromycin from pediatric suspension in infants and children. <i>Scandinavian journal of infectious diseases</i>. 1981 Sep 1;13(3):211-5.</p> <p>16. Murchison L, De Coppi P, Eaton S. Post-natal erythromycin exposure and risk of infantile hypertrophic pyloric stenosis: a systematic review and meta-analysis. <i>Pediatr Surg Int</i>. 2016;32:1147-52.</p> <p>17. Micromedex. Accessed online 29 May 2018.</p> <p>18. Curry JJ, Lander AD, Stringer MD, Committee BMR. A multicenter, randomized, double-blind, placebo-controlled trial of the prokinetic agent erythromycin in the postoperative recovery of infants with gastroschisis. <i>J Pediatr Surg</i>. 2004;39:565-9.</p> <p>19. Razzaq A, Safdar CA, Ali S. Erythromycin establishes early oral feeding in neonates operated for congenital intestinal atresias. <i>Pediatric Surgery International</i>. 2009;25:361-4.</p>
--	---

Original version Date: 20/06/2018	Author: NMF Consensus Group
Current Version number: 1.0	Current Version Date: 20/06/2018
Risk Rating: Low	Due for Review: 20/06/2023
Approval by: As per Local policy	Approval Date:

Authors Contribution

Original author/s	Chris Wake, Srinivas Bolisetty
Revision author/s	
Expert review	Tony Lai, Brendan McMullan
Evidence Review	David Osborn
Nursing Review	Eszter Jozsa, Robyn Richards
Pharmacy Review	Jing Xiao, Carmen Burman, Cindy Chen
NMF Group contributors	Nilkant Phad, Himanshu Popat, Michael Hewson, Jessica Ryan, Joanne Patel, Roland Broadbent
Final content and editing review of the original	Ian Whyte
Electronic version	Cindy Chen, Ian Callander
Facilitator	Srinivas Bolisetty