

<b>Indication</b>	<p>1. Prevention of iron-deficiency anaemia in infants at risk of reduced body stores</p> <ul style="list-style-type: none"> <li>-Preterm infants &lt;1800g</li> <li>-Infants 1800-2500g</li> <li>-Ex-preterm infants not tolerating feeds of 180 mL/kg/day with iron containing fortifier or formula</li> <li>-When ceasing iron containing fortifier or formula prior to discharge</li> </ul> <p>2. Treatment of iron deficiency anaemia</p> <p>NOTE: Formulas with 5-9 mg/L of iron are considered adequate in iron content</p> <p style="text-align: center;"><b>Iron content in common Fortifiers and Formulas at various mL/kg/day:</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th colspan="3" style="text-align: center;">Iron content</th> </tr> <tr> <th></th> <th style="text-align: center;">140 mL/kg/day</th> <th style="text-align: center;">160 mL/kg/day</th> <th style="text-align: center;">180 mL/kg/day</th> </tr> </thead> <tbody> <tr> <td><b>Preterm EBM</b></td> <td style="text-align: center;">0.04 mg/kg/day</td> <td style="text-align: center;">0.05 mg/kg/day</td> <td style="text-align: center;">0.054 mg/kg/day</td> </tr> <tr> <td><b>EBM+S26 HMF</b></td> <td style="text-align: center;">0.04 mg/kg/day</td> <td style="text-align: center;">0.05 mg/kg/day</td> <td style="text-align: center;">0.054 mg/kg/day</td> </tr> <tr> <td><b>EBM+FM 85</b></td> <td style="text-align: center;">2.1 mg/kg/day</td> <td style="text-align: center;">2.4 mg/kg/day</td> <td style="text-align: center;">2.7 mg/kg/day</td> </tr> <tr> <td><b>EBM+Nutricia BMF</b></td> <td style="text-align: center;">0.04 mg/kg/day</td> <td style="text-align: center;">0.05 mg/kg/day</td> <td style="text-align: center;">0.054 mg/kg/day</td> </tr> <tr> <td><b>Neocate Gold</b></td> <td style="text-align: center;">1.4 mg/kg/day</td> <td style="text-align: center;">1.6 mg/kg/day</td> <td style="text-align: center;">1.8 mg/kg/day</td> </tr> <tr> <td><b>Pre Nan Gold</b></td> <td style="text-align: center;">2.5 mg/kg/day</td> <td style="text-align: center;">2.9 mg/kg/day</td> <td style="text-align: center;">3.2 mg/kg/day</td> </tr> <tr> <td><b>Aptamil Gold + Preterm</b></td> <td style="text-align: center;">2.2 mg/kg/day</td> <td style="text-align: center;">2.6 mg/kg/day</td> <td style="text-align: center;">2.9 mg/kg/day</td> </tr> <tr> <td><b>S26LBW</b></td> <td style="text-align: center;">2.0 mg/kg/day</td> <td style="text-align: center;">2.2 mg/kg/day</td> <td style="text-align: center;">2.5 mg/kg/day</td> </tr> <tr> <td><b>Elecare/Elecare LCP</b></td> <td style="text-align: center;">1.7 mg/kg/day</td> <td style="text-align: center;">1.9 mg/kg/day</td> <td style="text-align: center;">2.2 mg/kg/day</td> </tr> <tr> <td><b>Pepti-Junior</b></td> <td style="text-align: center;">1 mg/kg/day</td> <td style="text-align: center;">1.2 mg/kg/day</td> <td style="text-align: center;">1.4 mg/kg/day</td> </tr> </tbody> </table>		Iron content				140 mL/kg/day	160 mL/kg/day	180 mL/kg/day	<b>Preterm EBM</b>	0.04 mg/kg/day	0.05 mg/kg/day	0.054 mg/kg/day	<b>EBM+S26 HMF</b>	0.04 mg/kg/day	0.05 mg/kg/day	0.054 mg/kg/day	<b>EBM+FM 85</b>	2.1 mg/kg/day	2.4 mg/kg/day	2.7 mg/kg/day	<b>EBM+Nutricia BMF</b>	0.04 mg/kg/day	0.05 mg/kg/day	0.054 mg/kg/day	<b>Neocate Gold</b>	1.4 mg/kg/day	1.6 mg/kg/day	1.8 mg/kg/day	<b>Pre Nan Gold</b>	2.5 mg/kg/day	2.9 mg/kg/day	3.2 mg/kg/day	<b>Aptamil Gold + Preterm</b>	2.2 mg/kg/day	2.6 mg/kg/day	2.9 mg/kg/day	<b>S26LBW</b>	2.0 mg/kg/day	2.2 mg/kg/day	2.5 mg/kg/day	<b>Elecare/Elecare LCP</b>	1.7 mg/kg/day	1.9 mg/kg/day	2.2 mg/kg/day	<b>Pepti-Junior</b>	1 mg/kg/day	1.2 mg/kg/day	1.4 mg/kg/day
	Iron content																																																
	140 mL/kg/day	160 mL/kg/day	180 mL/kg/day																																														
<b>Preterm EBM</b>	0.04 mg/kg/day	0.05 mg/kg/day	0.054 mg/kg/day																																														
<b>EBM+S26 HMF</b>	0.04 mg/kg/day	0.05 mg/kg/day	0.054 mg/kg/day																																														
<b>EBM+FM 85</b>	2.1 mg/kg/day	2.4 mg/kg/day	2.7 mg/kg/day																																														
<b>EBM+Nutricia BMF</b>	0.04 mg/kg/day	0.05 mg/kg/day	0.054 mg/kg/day																																														
<b>Neocate Gold</b>	1.4 mg/kg/day	1.6 mg/kg/day	1.8 mg/kg/day																																														
<b>Pre Nan Gold</b>	2.5 mg/kg/day	2.9 mg/kg/day	3.2 mg/kg/day																																														
<b>Aptamil Gold + Preterm</b>	2.2 mg/kg/day	2.6 mg/kg/day	2.9 mg/kg/day																																														
<b>S26LBW</b>	2.0 mg/kg/day	2.2 mg/kg/day	2.5 mg/kg/day																																														
<b>Elecare/Elecare LCP</b>	1.7 mg/kg/day	1.9 mg/kg/day	2.2 mg/kg/day																																														
<b>Pepti-Junior</b>	1 mg/kg/day	1.2 mg/kg/day	1.4 mg/kg/day																																														
<b>Action</b>	<p>Iron is needed for the production of haemoglobin and certain iron-containing enzymes. Ferrous sulphate corrects iron deficiency by re-saturating iron storage organs.</p>																																																
<b>Drug Type</b>	<p>Iron supplement</p>																																																
<b>Trade Name</b>	<p>Ferro-Liquid</p>																																																
<b>Dosage / Interval</b>	<p><b>Prophylaxis:</b> 2 mg/kg of <b>elemental iron</b> daily. To commence at 2-6 weeks of age (2-4 weeks of age in extremely low-birthweight infants)</p> <p><b>Treatment:</b> 2 mg/kg of <b>elemental iron</b> 12 hourly</p>																																																
<b>Maximum daily dose</b>	<p>15 mg daily INDEPENDENT OF WEIGHT</p> <p>Prophylaxis: 2-3 mg/kg/day of <b>elemental iron</b>. Commence only when on full feeds in hospital or prior to discharge when fortification is stopped or preterm formula changed to term formula. Delay in infants with multiple transfusions and increased Serum Ferritin levels (&gt;350 microgram/L) or have received a transfusion in last 2 weeks. Prophylaxis dose &gt;5 mg/kg/day should be avoided in preterm infants because of possible risk of retinopathy of prematurity</p> <p>Treatment: Can commence on 3 mg/kg/day of <b>elemental iron</b> and may need to go up to 6 mg/kg/day in iron deficiency anaemia or on erythropoietin. It is suggested to undertake iron studies to titrate the dose.</p>																																																
<b>Total cumulative dose</b>	<p>Doses &gt;5 mg/kg/day should be avoided in preterm infants because of possible risk of retinopathy of prematurity</p>																																																
<b>Presentation</b>	<p>Ferrous sulphate liquid: 150 mg <b>ferrous sulfate</b>/5 mL equivalent to 30 mg <b>elemental iron</b>/5 mL=6 mg/mL of <b>elemental iron</b></p>																																																
<b>Route</b>	<p>Oral</p>																																																
<b>Administration</b>	<p>Oral or intragastric tube</p>																																																
<b>Monitoring</b>	<p>Periodic haemoglobin and reticulocyte count during therapy. Can take 2 weeks for haemoglobin concentrations to rise.</p> <p>Regular serum ferritin if treating iron deficiency anaemia.</p>																																																
<b>Contraindications</b>	<p>Haemolytic anaemia, haemochromatosis, haemosiderosis</p>																																																

<b>Precautions</b>	Excessive iron supplementation can lead to increased risk of infection, poor growth and disturbed absorption or metabolism of other minerals. Being a potent pro oxidant, non-protein bound iron can cause free oxygen radicals and increase risk of retinopathy of prematurity, especially when given in high doses as a component of blood transfusions or as adjunct to erythropoietin therapy. Risk of iron induced haemolysis in preterm infants with Vitamin E deficiency is more in first 6 weeks.
<b>Drug interaction</b>	Zinc supplementation does not impede iron absorption. There is no effect of iron supplementation on zinc or selenium absorption. Iron absorption from fortified milk is intact in spite of its high calcium content.
<b>Adverse Reactions</b>	GI irritation: epigastric pain, diarrhoea, constipation, dark stools (green or black), erosion of gastric mucosa Increased RBC haemolysis and haemolytic anaemia in premature infants with low vitamin E levels Rickets - with large doses of iron over a prolonged period of time. Acute toxicity - more severe GI effects including haematemesis and malaena, lethargy, pallor, cyanosis and shock
<b>Compatibility</b>	Can be administered with Penta-vite.
<b>Stability</b>	Solution may be used up to one month after opening (document date of opening on label).
<b>Storage</b>	Store below 25°C. Protect from light
<b>Special comments</b>	Infants on erythropoietin or infants with uncompensated blood loss may initially need higher doses and could be receiving iron supplementation in addition to preterm formula or fortified human milk.
<b>Evidence summary</b>	<p>1. LBW infant supplementation-RCT (Berglund 2010) suggested all babies born less than 2.5 kg will benefit from early supplementation, main outcome being iron deficiency at 6 months</p> <p>2.Dose - Low vs high-RCT (Friel 2001)-Preterm infants with an average birth weight of 1.46 kg received an iron intake of 5.9 versus 3.0 mg per kg per day at discharge and about 3 versus 2mg per kg per day at 3 to 9 months. There was no difference between the 2 groups in anaemia prevalence or neurodevelopment at 12 months, but the high-iron group had higher glutathione peroxidase concentrations (a marker of oxidative stress), lower plasma zinc and copper levels, and more respiratory tract infections, suggesting possible adverse effects from the higher intake.</p> <p>3.Early vs late supplementation-3 RCTs ( Franz 2000,Stienmarcher 2007, Arnon 2009)</p> <p>Steinmacher 2007-showed increased proportion of children with abnormal clinical neurological examination in late iron group (35% vs 17%;p=0.02)</p> <p>Franz et al randomised 204 infants with an average birth weight of 0.87 kg into an early iron group receiving 2 to 4 mg/kg/day of iron supplements from about 2 weeks and a late iron group that did not receive iron supplements until 2 months of age. There were no differences in serum ferritin and haematocrit at 2 months of age but infants in the late iron group had received more blood transfusions.</p> <p>Arnon 2009 - large RCT, 2 weeks vs 4 weeks, improved dermatological parameters in early group.</p>
<b>References</b>	<p>1.Hey E, Neonatal Formulary 4th edition. London: BMJ Publishing Group, 2003, pp87-88</p> <p>2.Neonatal and Paediatric Pharmacists Group, Royal College of Paediatrics and Child Health: Medicines for Children 2nd Edition, London: RCPCH Publications Limited, 2003, pp343-345</p> <p>3.Young TE, Mangum B: Neofax: A Manual of Drugs Used in Neonatal Care, 13th edition. Raleigh, North Carolina: Acorn Publishing, USA, 2000, pp 198</p> <p>4.Ferro-Liquid: Product Information (MIMSONline)</p> <p>5.Agostoni C, Buonocore G, Carnielli VP et al for the ESPGHAN Committee on Nutrition. Enteral nutrient supply for preterm infants: Commentary from the European Society for Paediatric Gastroenterology, Hepatology and Nutrition Committee on Nutrition. Journal of Pediatric Gastroenterology and Nutrition 2010;50:1-9</p> <p>6.Neofax 2011. Twenty fourth edition. Thomson Reuters, Montvale, NJ.</p> <p>7.Franz AR, Mihatsch WA, Sander S, Kron M, Pohlandt F. Prospective randomised trial of early vs late enteral iron supplementation in infants with a birth weight less than 1301grams. Pediatrics 2000;106:700-706</p> <p>8.Sankar MJ, Saxena R, Mani K et al. Early iron supplementation in very low birth weight infants – a randomised trial. Acta</p>

	<p>Paediatrica 2009;96:953-58</p> <p>9. Berseth CL, Van Aerde JE, Gross S, et al. Growth, efficacy in safety of feeding an iron fortified human milk fortifier. Pediatrics 2004;111:e699-e706.</p> <p>10. Berglund S, Westrup B, Domellof M. Iron supplements reduce the risk of iron deficiency anemia in marginally low birth weight infants. Pediatrics. 2010; 126(4): e874-83.</p> <p>11. Young L, Embleton ND, McCormick FM, McGuire W. Multinutrient fortification of human breast milk for preterm infants following hospital discharge. The Cochrane database of systematic reviews. 2013;2:CD004866.</p> <p>12. Rao R, Georgieff MK. Iron therapy for preterm infants. Clinics in Perinatology 2009;36:27-42</p> <p>13. Mills RJ, Davies MW. Enteral iron supplementation in preterm and low birth weight infants. The Cochrane database of systematic reviews. 2012</p> <p>14. American Academy of Pediatrics, Committee on Nutrition. Nutritional needs of the preterm infant. In: Kleinman, RE., editor. Pediatric nutrition handbook. Vol. 5. Chapel Hill, NC: American Academy of Pediatrics; 2004. p. 23-54.</p> <p>15. Carbonell-Estrany X, Figueras-Aloy J, Alvarez E. Erythropoietin and prematurity--where do we stand? J Perinat Med 2005;33(4):277-286.</p> <p>16. Barclay SM, Lloyd DJ, Duffy P, Aggett PJ. Iron supplements for preterm or low birthweight infants. Arch Dis Child Nov;1989 64(11):1621-1622.</p> <p>17. Halliday HL, Lappin TR, McClure G. Iron status of the preterm infant during the first year of life. Biol Neonate 1984;45(5):228-235.</p> <p>18. Steinmacher J, Pohlandt F, Bode H, Sander S, Kron M, Franz AR. Randomized trial of early versus late enteral iron supplementation in infants with a birth weight of less than 1301 grams: neurocognitive development at 5.3 years' corrected age. Pediatrics Sep;2007 120(3):538-54</p> <p>19. Arnon S, Regev RH, Bauer S, Shainkin-Kestenbaum R, Shiff Y, Bental Y, et al. Vitamin E levels during early iron supplementation in preterm infants. American Journal of Perinatology 2009;26:387-92.</p> <p>20. Barclay SM, Aggett PJ, Lloyd DJ, Duffy P. Reduced erythrocyte superoxide dismutase activity in low birth weight infants given iron supplements. Pediatric Research 1991;29:297-301. Andrews WL, Friel JK, Simmons B, Aziz K, Kwa PG, Downton G. Effect of two levels of iron supplementation of infant formulas on developmental outcome and iron nutrition in very low birth weight infants. Pediatric Research 1998;43:255A.</p> <p>23. Friel JK, Andrews WL, Aziz K, Kwa PG, Lepage G, L'Abbe MR. A randomized trial of two levels of iron supplementation and developmental outcome in low birth weight infants. Journal of Pediatrics 2001;139:254-60.</p>
--	--

<b>Original version Date: 08/08/2015</b>	<b>Author: NeoMed Consensus Group</b>
<b>Current Version number: 1</b>	<b>Version Date: 08/08/2015</b>
<b>Risk Rating: Medium</b>	<b>Due for Review: 08/08/2018</b>
<b>Approval by: As per Local policy</b>	<b>Approval Date: As per Local policy</b>